

6xLaemmli SDS sample Buffer

Catalog No.	Description	Size
SAB03-01	6x Laemmli SDS sample Buffer	25 ml
SAB03-02	6x Laemmli SDS sample Buffer	50 ml

Description:

The Laemmli SDS sample Buffer is used for the preparation of protein samples for SDS-PAGE. The buffer contains all the necessary reagents for complete disruption of high-order protein structures. The SDS included in the buffer binds to hydrophobic regions of the protein, causing the protein to unfold and giving it a negative charge. The reducing agent (β -mercaptoethanol or DTT) breaks disulfide bonds and destroys residual secondary structures. As β -mercaptoethanol or DTT is prone to oxidation during multiple freeze-thaw cycles, it needs to be added before use.

Composition

Tris HCl (pH 6.8)	375 mM
SDS	9%
Glycerol	50%
Bromophenol blue	0.03%

Note: Before use, add 90 μ l β -mercaptoethanol is needed for 910 μ l 6x Laemmli SDS Sample Buffer.

Storage Temperature: Ambient temperature

Procedure

1. Dissolve precipitated solids (if any) in the Sample Buffer at 37°C.
2. Mix gently the Sample Buffer to ensure that the solutions are homogeneous.
3. Add 90 μ l β -mercaptoethanol to 910 μ l 6x Laemmli SDS sample Buffer and mix.
4. Add **1 μ l** of 6x sample Buffer to **5 μ l** protein sample and mix. Using 1x Sample Buffer (dilute in ddH₂O or Milli-Q H₂O) to make up the volume difference if needed.
5. Heat samples at 100°C for 5 minutes.
6. Centrifuge briefly and load directly to a SDS polyacrylamide gel.



Bioland Scientific LLC 14925 Paramount Blvd., Suite C, Paramount, CA 90723

Tel: (562)602-8882 Fax: (562)733-6008 Email: service@bioland-sci.com Online: www.bioland-sci.com